Edinburgh Postnatal Depression Scale (EPDS)

Date:	Clinic Name/Nur	nber:		
Your Age:	Weeks o	f Pregnancy/	'Age of Baby:	
the blank by the answer 10 items and find your	that comes closest to how score by adding each number	you have felt r that appear	IN THE PAST 7 DAYS—not just in parentheses (#) by your cl	ne place a CHECK MARK () on st how you feel today. Complete all necked answer. This is a provider regardless of your score.
Below is an example alı	ready completed.		7. I have been so unhappy sleeping:	that I have had difficulty
I have felt happy: Yes, all of the time Yes, most of the time No, not very often No, not at all		(0) (1) (2) (3)	Yes, most of the time Yes, sometimes No, not very often No, not at all 8. I have felt sad or misera	
	ave felt happy most of the tin complete the other question		Yes, most of the time Yes, quite often Not very often No, not at all	(3) (2) (1) (0)
 I have been able to things: As much as I alway: Not quite so much is Definitely not so much is Not at all 	now	(0) (1) (2) (3)	9. I have been so unhappy Yes, most of the time Yes, quite often Only occasionally No, never	
2. I have looked forwa As much as I ever of Rather less than I u Definitely less than Hardly at all	used to	(0) (1) (2) (3)	10. The thought of harming Yes, quite often Sometimes Hardly ever Never	myself has occurred to me:* (3)(2)(1)(0)
·			* If you scored a 1, 2 or 3 or HEALTH CARE PROVIDER (OB midwife) OR GO TO THE EMEL own safety and that of your ball your total score is 11 or man postpartum depression (PPD)	RGENCY ROOM NOW to ensure your by. ore, you could be experiencing or anxiety. PLEASE CALL YOUR
4. I have been anxious No, not at all Hardly ever Yes, sometimes Yes, very often	s or worried for no good reas	on: (0) (1) (2) (3)	week or call your health care nurse-midwife).	
5. I have felt scared o Yes, quite a lot Yes, sometimes No, not much No, not at all	r panicky for no good reason	:(3)(2)(1)(0)	that make them cry or get ang normal. However, if they worse	ry easily. Your feelings may be n or continue for more than a week ovider (OB/Gyn, family doctor or r can be a new and stressful elf by:
cope at all Yes, sometimes I ha as usual	ne I haven't been able to aven't been coping as well be I have coped quite well	(3) (2) (1) (0)	or anxiety, please contact your	t's just walking outside. I have concerns about depression health care provider. epression Scale (EPDS) is a screening tool
			_	

Edinburgh Postnatal Depression Scale (EPDS) Scoring & Other Information

ABOUT THE EPDS

Studies show that postpartum depression (PPD) affects at least 10 percent of women and that many depressed mothers do not get proper treatment. These mothers might cope with their baby and with household tasks, but their enjoyment of life is seriously affected, and it is possible that there are long term effects on the family.

The Edinburgh Postnatal Depression Scale (EPDS) was developed to assist health professionals in detecting mothers suffering from PPD; a distressing disorder more prolonged than the "blues" (which can occur in the first week after delivery).

The scale consists of 10 short statements. A mother checks off one of four possible answers that is closest to how she has felt during the past week. Most mothers easily complete the scale in less than five minutes.

Responses are scored 0, 1, 2 and 3 based on the seriousness of the symptom. Items 3, 5 to 10 are reverse scored (i.e., 3, 2, 1, and 0). The total score is found by adding together the scores for each of the 10 items.

Mothers scoring above 12 or 13 are likely to be suffering from depression and should seek medical attention. A careful clinical evaluation by a health care professional is needed to confirm a diagnosis and establish a treatment plan. The scale indicates how the mother felt during the previous week, and it may be useful to repeat the scale after two weeks.

INSTRUCTIONS FOR USERS

- 1. The mother checks off the response that comes closest to how she has felt during the previous seven days.
- 2. All 10 items must be completed.
- 3. Care should be taken to avoid the possibility of the mother discussing her answers with others.
- 4. The mother should complete the scale herself, unless she has limited English or reading difficulties.
- 5. The scale can be used at six to eight weeks after birth or during pregnancy.

Please note: Users may reproduce this scale without further permission providing they respect the copyright (which remains with the British Journal of Psychiatry), quote the names of the authors and include the title and the source of the paper in all reproduced copies. Cox, J.L., Holden, J.M. and Sagovsky, R. (1987). Detection of postnatal depression: Development of the 10-item Edinburgh Postnatal Depression Scale. British Journal of Psychiatry, 150, 782-786.